

## Philadel studied the Fourier coefficients

$$h(q) = 1 + \sum_{n=1}^{\infty} \frac{q^{n(n+1)/2}}{(1+q)(1+q^2)\cdots(1+q^n)};$$

where  $h$  is a function that appears in the work of Ramanujan. They prove, among other things, that, for  $m > 0$  and  $m \equiv 1 \pmod{24}$ , that these Fourier coefficients are given by

$$T(m) = \# \left( \begin{array}{l} \text{equivalence classes } [(x; y)] \text{ of solutions to} \\ x^2 - 6y^2 = m \text{ with } x + 3y \equiv 1 \pmod{12} \end{array} \right) \\ = \# \left( \begin{array}{l} \text{equivalence classes } [(x; y)] \text{ of solutions to} \\ x^2 - 6y^2 = m \text{ with } x + 3y \equiv 5 \pmod{12} \end{array} \right);$$

Cohen showed that

$$h_0(\tau) = y^{1/2} \sum_{\substack{n \in \mathbb{Z} \\ n \neq 0}} T(n) e^{inx=24} K_0 \frac{2-j\eta y}{24}$$

is a Maass waveform on  $h_0(2)$ . Zweger was able to place  $h_0(\tau)$  in a larger framework of indefinite theta functions.

In this talk, I will discuss the problem of placing quadratic identities arising in the work of ADH into a modular framework. This is joint work, in progress, with Larry Rolin.

Wednesday, February 21, 2018, 2:40 { 4:00 PM

Bryn Mawr College, Department of Mathematics

Park Science Center 328 Tea and refreshments at 2:20PM in Park 339